

ULTRA-PERIPHERAL COLLISIONS IN STAR

P. YEPES

for the STAR Collaboration

Rice University

MS 315, 6100 Main Street, Houston, TX 77005

E-mail: yepes@rice.edu

Ultra-peripheral heavy ion collisions involve long range electromagnetic interactions at impact parameters larger than twice the nuclear radius, where no hadronic nucleon-nucleon collisions occur. The first observation of coherent ρ^0 production with and without accompanying nuclear breakup, along with the observation of e^+e^- pair production are reported by the STAR collaboration.

1 Introduction

We refer to ultra-peripheral heavy ion collisions, as those nuclear interactions with a impact parameter, b , larger than twice the nuclear radius R_A , where no hadronic nucleon-nucleon collisions occur¹. The large charge of a relativistic heavy nucleus is a strong source of quasi-real photons. Exclusive ρ^0 meson production $AuAu \rightarrow AuAu\rho^0$ can then be described by the vector meson dominance model²: a photon emitted by one nucleus fluctuates to a virtual quark-anti quark pair; this intermediate state scatters diffractively from the other nucleus, emerging as a vector meson. The diagram for this process is shown in Fig. 1a. Here, the gold nuclei remain in their ground state. Additional photon exchange can yield nuclear excitation and the subsequent emission of single or multiple neutrons, as shown in Fig. 1b for the process $AuAu \rightarrow Au^*Au^*\rho^0$.

Photon and Pomeron can couple coherently to the spatially extended




Figure 1. Feynman diagrams for (a) ρ^0 production, and (b) ρ^0 production with nuclear excitation. Factorization is indicated by the dashed line; nuclear excitation may occur by the exchange of a single or multiple photons.

electric and nuclear charge of the gold nuclei. Requiring a coherent interaction of the whole nucleus imposes a maximum momentum transfer of $\hbar/d$, where d is the nuclear dimension. In the transverse direction along which nuclei are not Lorentz contracted, d is of the order of the nucleus radius, and therefore $p_T < \hbar/R_A$ (~ 100 MeV/c for $R_A \sim 7$ fm). In the longitudinal direction nuclei are contracted by a factor γ , therefore the maximum longitudinal momentum is γ times larger: $p_{\parallel} < \hbar\gamma/R_A$ (~ 6 GeV/c at RHIC).

The coupling strength of the photon is proportional to the square of the charge Z^2 (6241 for $AuAu$); the strength of the Pomeron coupling lies between $A^{4/3}$ for surface coupling to A^2 in the bulk limit (10^3 to 10^4 for $AuAu$). For gold collisions at $\sqrt{s_{NN}} = 130$ GeV those large couplings translate into a ρ production cross section of about 400 mb, or $\approx 5\%$ of the total hadronic cross section.

In the case of ρ production, it is not possible to determine which nucleus is the photon source and which is the target. Therefore the amplitudes for ρ^0 production from both ions interfere. Since the ρ^0 has negative parity, this interference is destructive. The short-lived ρ^0 decay before they travel the distance of the impact parameter b , and the interference is believed to be sensitive to the post-decay wave function⁶.




Figure 2. End and side views of a typical ρ candidate event in the STAR TPC. The candidate tracks are almost back-to-back radially, but boosted longitudinally.

The ρ meson decays into $\pi\pi$ with an approximately 100% branching ratio. However a $\pi\pi$ final state may be produced directly or through the ρ . The amplitudes for ρ production, A , and direct $\pi\pi$ production, B , interfere

and⁴

$$\frac{d\sigma}{dM_{\pi\pi}} = \left| \frac{A\sqrt{M_{\pi\pi}M_\rho\Gamma_\rho}}{M_{\pi\pi}^2 - M_\rho^2 + iM_\rho\Gamma_\rho} + B \right|^2 \quad (1)$$

where the ρ width is corrected for the increasing phase space as $m_{\pi\pi}$ increases. The ρ component undergoes a 180° phase shift at M_ρ , so the interference skews the ρ peak shape, enhancing production for $M_{\pi\pi} < M_\rho$ and suppressing the spectrum for $M_{\pi\pi} > M_\rho$.

2 STAR Detector

In the year 2000, RHIC collided gold nuclei at a center-of-mass energy of $\sqrt{s_{NN}} = 130$ GeV/nucleon. The STAR detector consists of a 4.2 m long cylindrical time projection chamber (TPC) of 2 m radius. In 2000 the TPC was operated in a 0.25 T solenoidal magnetic field. Particles are identified by their energy loss in the TPC. A central trigger barrel of scintillators surrounds the TPC, providing fast signaling for online event selection. Two zero degree calorimeters (ZDC) at ± 18 m from the interaction point are sensitive to the neutral remnants of nuclear breakup.

3 Analysis

Production of the ρ^0 meson in ultra-peripheral collisions has a clear experimental signature, especially when compared with the typical high multiplicity AuAu interaction. It is characterized by a $\pi^+\pi^-$ pair in an otherwise 'empty' spectrometer. Fig. 1 shows a typical event candidate; the tracks are approximately back-to-back in the transverse plane due to the small p_T of the pair. Two data sets are used for the analysis: the 'minimum bias' and the 'topology' samples.

3.1 Minimum Bias Sample

The former includes 800k events triggered by requiring simultaneous signal in both ZDC's. The analysis of the 'minimum bias' sample, on which this paper focuses, is optimized to select the following reaction: $AuAu \rightarrow Au^*Au^*\rho^0$. Events are required to have two tracks of opposite charge coming from a common vertex. In addition the vertex should be within 15cm from the beam axis, and contained in the TPC in the longitudinal direction. Moreover the opening angle in the transverse plane is required to be larger than 2.7 radians.

A sample of 100K Monte Carlo $AuAu \rightarrow AuAu\rho^0$ events, generated with STARLIGHT⁵, and run through the STAR simulation and reconstruction




Figure 3. Reconstruction efficiency, including acceptance, as a function of the rapidity, for the inclusive ρ^0 production.




Figure 4. ρ^0 meson reconstruction efficiency, including acceptance, as a function of transverse momentum, before the cut in the opening angle is applied. Only ρ^0 mesons with $|\eta| < 1$ were considered.

chain was utilized to estimate the reconstruction efficiency for ρ^0 inclusive production. The efficiency, including geometrical acceptance, as a function of the ρ^0 rapidity is depicted in Fig. 3. As can be seen the efficiency peaks at 70% at mid rapidity. The dip at rapidity zero is due to the poor vertex resolution for those events. This is due to the fact that it is not possible to determine a common vertex for two tracks perfectly back-to-back. Fig. 4 depicts the efficiency as a function of transverse momentum, before the cut on the angle in the transverse plane is applied. That cut affects the high end of the distribution, but has little effect in the overall acceptance, since ρ^0 p_T distribution peaks below 100 MeV/c.

Fig. 5.a shows the uncorrected transverse momentum spectrum for the selected events, along with estimated background from hadronic processes and dielectron events. A clear peak, the signature for coherent coupling, can be observed at $p_T < 100$ MeV/c. Those events are compatible with coherently produced ρ^0 candidates. The hadronic background is estimated from like-sign combination pairs, which is normalized to the signal for $p_T > 200$ MeV/c. As can be noted, the hadronic background does not have a peak at low p_T . Di-




Figure 5. a) The uncorrected p_T spectrum for ρ^0 candidates in the minimum bias sample, along with estimates for the hadronic and di-electron backgrounds. b) The p_T spectrum for ρ^0 candidates after background subtraction and correcting for acceptance, reconstruction efficiency, and smearing effects.

electron background is estimated by selecting electrons in the kinematic zone where the energy loss dE/dx in the TPC provides good separation between electron and pion as explained below. Fig. 5.b shows the corrected transverse momentum after background subtraction, and correction for acceptance, detection efficiency, and smearing effects.

Fig. 6.a depicts the $\pi\pi$ invariant mass distribution for the events with a total p_T smaller than 150 MeV/c, along with estimates for hadronic and di-electron backgrounds. As can be observed, a clear peak is shown at the ρ^0 mass, while backgrounds are concentrated at low masses. The spectrum corrected for background, acceptance, reconstruction efficiency and smearing is shown in Fig. 6.b. The corrected spectrum is well fit by Eq. 1, with $|B/A|$ similar to that found by the ZEUS collaboration for $\gamma p \rightarrow \rho^0 p^7$.




Figure 6. a) The uncorrected $m_{\pi\pi}$ spectrum of ρ^0 candidates with $p_T < 150$ MeV/c, along with estimates for hadronic and di-electron backgrounds. b) The corrected $m_{\pi\pi}$ spectrum of ρ^0 candidates with $p_T < 150$ MeV/c along with the results from a fit to equation(1).

3.2 Topology Sample

For the analysis of the process $AuAu \rightarrow AuAu\rho^0$ a data sample selected with a low-multiplicity topology trigger was utilized. In the level 0 trigger, the central trigger barrel was divided into 16 coarse pixels. These pixels cover half a unit in rapidity, and 90° in the transverse plane. Hits in opposite pixels were required, while pixels in the top and the bottom acted as a veto to suppress cosmic rays. No requirement on the ZDC signals was imposed. A high level trigger⁸ further removed background. The STAR collaboration collected about 30k events in 7 hours, with this configuration. The ρ^0 candidates from this data set have a transverse momentum and an invariant mass distributions similar to the ones already shown in Figures 5 and 6. Therefore, they are also characterized by a p_T distribution peaked below 100 MeV/c, and an events, around 300, clustered around the ρ^0 mass. In contrast to the minimum bias data, the topology triggered data had almost no energy deposition in the ZDC

consistent with the two gold nuclei remaining in their ground state.

3.3 Electron-Positron pair production

Two-photon interactions include the purely electromagnetic process of electron-positron pair production as well as single and multiple meson production. The coupling $Z\alpha$ (0.6 for Au) is large, hence e^+e^- pair production is an important probe of quantum electrodynamics in strong fields¹. At momenta below 140 MeV/c, e^+e^- pairs are identified by their energy loss in the TPC as shown for the minimum bias data sample in Fig. 7a. Fig. 7b shows the p_T spectrum for identified e^+e^- pairs; a clear peak at $p_T < 50$ MeV/c identifies the process $AuAu \rightarrow Au^*Au^*e^+e^-$.




Figure 7. (a) Energy loss dE/dx of tracks in the 2-track, minimum bias data; triangles indicate events where both particles are identified as electrons. (b) The p_T spectrum for identified e^+e^- pairs.

4 Summary

In summary, for the first time, exclusive ρ^0 production $AuAu \rightarrow AuAu\rho^0$ and ρ^0 production with and without nuclear breakup was observed in ultra-peripheral heavy ion collisions. The ρ^0 are produced at small perpendicular momentum, showing their coherent coupling to both nuclei. In addition, the coherent electromagnetic process $AuAu \rightarrow Au^*Au^*e^+e^-$ was observed.

References

1. G. Baur, K. Hencken and D. Trautmann, J. Phys. G**24**, 1657 (1998); C. A. Bertulani and G. Baur, Phys. Rep. **163**, 299 (1988).
2. J.J. Sakurai, Ann. Phys. **11** (1960) 1, and Phys. Rev. Lett. **22** 981 (1969), T.H. Bauer *et al.*, Rev. Mod. Phys. **50** 261 (1978).
3. S. Klein and J. Nystrand, Phys. Rev. **C60**, 014903 (1999).

4. P. Söding, Phys. Lett. **19**, 702 (1966).
5. S. Klein and J. Nystrand, private communication.
6. S. Klein and J. Nystrand, Phys. Rev. Lett. **84**, 2330 (2000).
7. J. Breitweg *et al.*, Eur. Phys. J **C2**, 247 (1998).
8. J. S. Lange *et al.*, Nucl. Instrum. Meth. **A453**, 397 (2000).